

# BULLETIN BOARD

## LETTERS

### Pharmacists and POEC

Madam

We read with interest and some surprise the description of the 3-day training course provided in Lambeth, Southwark and Lewisham, London<sup>1</sup> for pharmacists, enabling them to issue progestogen-only emergency contraception (POEC) using a Patient Group Direction (PGD).

First, do they need that much training? As 'very streetwise' professionals, do they need so much training to issue a drug, when the World Health Organization (WHO) *Medical Eligibility Criteria for Contraceptive Use*<sup>2</sup> advise that there are no medical contraindications to POEC and the Faculty's own guidelines also state that there are no absolute contraindications to POEC, although caution should be used in women with porphyria or severe liver disease.<sup>3</sup> Second, how on earth did they find the time?

We, in Worcestershire, train pharmacists over two evenings: the first evening for pharmacists entering the scheme; the second evening for pharmacists experienced in issuing emergency contraception under PGDs. There is a sharing of experience, likes and dislikes about offering the service, how to train shop staff to be supportive, dealing with press enquiries, revisiting child protection issues, etc.

The sessions were deemed to be valuable, informative, fun and useful and, with a 24-hour sexual health consultant on-call rota, and several young people's outreach health services throughout the county, to whom they can refer, the pharmacists here feel fully supported and valued.

We believe barriers should not be created to women accessing emergency contraception – particularly the high price of over-the-counter (OTC) products. Decreasing the cost of OTC Levonelle<sup>®</sup> to that of a prescription would increase accessibility and sales. We should like to see school nurses trained and able to issue POEC and for easier access for all women who need this method of contraception.

**Yvonne Stedman**, MB BS, MFFP

*Joint Clinical Director, Contraception and Reproductive Healthcare, South Worcestershire NHS PCT, Sexual Health Service, Moor Street Clinic, 7 Moor Street, Worcester WR1 3DB, UK*

**Jenny Kimberlee**, Grad IPM, PgDip Health Education  
*Teenage Pregnancy Co-ordinator, Sexual Health Service, Moor Street Clinic, Worcester, UK*

#### References

- 1 Bacon L, Savage I, Cook S, Taylor B. Training and supporting pharmacists to supply progestogen-only emergency contraception. *J Fam Plann Reprod Health Care* 2003; **29**(2): 17–22.
- 2 Faculty of Family Planning and Reproductive Health Care Clinical Effectiveness Unit. FFRHC Guidance: Emergency Contraception (April 2003). *J Fam Plann Reprod Health Care* 2003; **29**(2): 9–16.
- 3 World Health Organization (WHO). Improving access to quality care in family planning. *Medical eligibility criteria for contraceptive use*. Geneva, Switzerland: WHO, 2000.

### Reply

We are grateful for the interest shown in our training.

After the appraisal of the first course, we did reduce the course to a 2-day one, as is stated in the discussion on page 21. Funding for locums was from the local Health Action Zone, although finding locums was a problem for the pharmacists concerned.

The main point to note is that the first course was held in early 2000, when the idea of pharmacists doing this work was very new (we were only the second project in the UK to go live) and when over-the-counter sale had not yet been approved. Public and professional reaction was untested, and a great deal of time was spent helping the pharmacists to feel confident if challenged about their right to supply emergency hormonal contraception. The wisdom of this was shown when the *Daily Mail* published an inaccurate story, as detailed in the paper; the pharmacists concerned had an extremely unpleasant experience but coped amazingly well. We were also anxious that the participants should see themselves as part of a seamless service including all sexual health providers, and had no idea of what demand might be (it has in fact reached 10 000 supplies a year across our very deprived inner city area).

Perhaps the moral of this is that today's daring innovation is tomorrow's boring received wisdom, as every department that has set up nurses supplying to a Patient Group Direction will recognise!

**Lesley Bacon**, MFFP, MRCGP

*Consultant in Sexual and Reproductive Health, Lewisham NHS PCT, Department of Reproductive Health, St Giles, St Giles Road, London SE5 7RN, UK*

**Beth Taylor**, BSc, MRPharmS

*Manager Community Services Pharmacy Team, Southwark NHS PCT, London, UK*

### Ectopic pregnancy following use of progestin-only ECPs

Madam

We read with great interest the editorial about ectopic pregnancy (EP) following use of progestin-only emergency contraceptive pills (ECPs).<sup>1</sup> The authors note that 12 EPs in women who used levonorgestrel ECPs have been reported in the UK and that a handful of additional cases have been reported in other countries. As the authors acknowledge, this information cannot be used to calculate the probability that a pregnancy occurring after use of the treatment will be ectopic because the total number of pregnancies needed for the denominator of the calculation is unknown. Nevertheless, based on these case reports, Britain's Committee on Safety of Medicines (CSM) has advised that if a woman who has used progestin-only ECPs becomes pregnant, "the possibility of an ectopic pregnancy should be considered".<sup>2</sup>

Data from clinical trials of ECPs can yield an accurate estimate of the rate of EP because pregnancies in these trials are systematically documented and thus provide a valid denominator for the rate. Through a search of

the published literature, we identified five clinical trials of levonorgestrel-only ECPs.<sup>3–7</sup> As shown in Table 1, these trials reported a total of 97 intrauterine pregnancies and one EP. The proportion of pregnancies that were ectopic was thus 1.02% (95% exact CI 0.02%–5.55%).

This proportion is consistent with the reported national rate of 12.4 and 19.7 per 1000 pregnancies in England and Wales and in the USA, respectively.<sup>8,9</sup> Therefore, these trials provide no evidence to suggest that progestin-only ECPs increase the chance that a pregnancy will be ectopic. Moreover, because ECPs are so effective at preventing pregnancy in general, they certainly reduce a woman's absolute risk of EP.

We agree with the authors of the Editorial and with the CSM that a woman who has used progestin-only ECPs and who subsequently has clinical symptoms of EP should seek appropriate evaluation, as should any woman with such symptoms. However, no evidence exists to warrant heightened concern in users of progestin-only ECPs.

**James Trussell**, PhD

*Professor of Economics and Public Affairs and Director, Office of Population Research, Wallace Hall, Princeton University, Princeton, NJ 08544, USA*

**Allison Hedley**, PhD

*Research Assistant, Office of Population Research, Princeton University, Princeton, NJ, USA*

**Elizabeth Raymond**

*Associate Medical Director, Clinical Trials Division, Family Health International, Research Triangle Park, NC, USA*

#### References

- 1 Harrison-Woolrych M, Woolley J. Progestogen-only emergency contraception and ectopic pregnancy. *J Fam Plann Reprod Health Care* 2003; **29**: 5–6.
- 2 Woolley J. Levonelle/Levonelle-2 emergency contraception: new advice. *CMO's Update* 35, January 2003: 9.
- 3 von Hertzen H, Piaggio G, Ding J, et al. Low dose mifepristone and two regimens of levonorgestrel for emergency contraception: a WHO multicentre randomized trial. *Lancet* 2002; **360**: 1803–1810.
- 4 Arowojolu AO, Okewole IA, Adekunle AO. Comparative evaluation of the effectiveness and safety of two regimens of levonorgestrel for emergency contraception in Nigerians. *Contraception* 2002; **66**: 269–273.
- 5 Task Force on Post-ovulatory Methods of Fertility Regulation. Randomised controlled trial of levonorgestrel versus the Yuzpe regimen of combined oral contraceptives for emergency contraception. *Lancet* 1998; **352**: 428–433.
- 6 Wu S, Wang C, Wang Y. A randomized, double-blind, multicentre study on comparing levonorgestrel and mifepristone for emergency contraception [in Chinese]. *Zhonghua Fu Chan Ke Za Zhi* 1999; **34**: 327–330.
- 7 Ho PC, Kwan MSW. A prospective randomized comparison of levonorgestrel with the Yuzpe regimen in post-coital contraception. *Hum Reprod* 1993; **8**: 389–392.
- 8 Rajkhowa M, Glass MR, Rutherford AJ, et al. Trends in the incidence of ectopic pregnancy in England and Wales from 1966 to 1996. *Br J Obstet Gynaecol* 2000; **107**: 369–374.
- 9 Ectopic pregnancies – United States, 1990 to 1992. *Morb Mortal Wkly Rep* 1995; **44**: 46–48.

**Table 1** Results of five clinical trials of levonorgestrel-only emergency contraceptive pills

| Trial                                | Known pregnancies (n) | Ectopic pregnancies (n) | Dose of levonorgestrel              |
|--------------------------------------|-----------------------|-------------------------|-------------------------------------|
| WHO (2002) <sup>3</sup>              | 1356                  | 24                      | 0.75 mg in two doses 12 hours apart |
| WHO (2002) <sup>3</sup>              | 1356                  | 20                      | 1.5 mg as a single dose             |
| Arowojolu et al. (2002) <sup>4</sup> | 545                   | 7                       | 0.75 mg in two doses 12 hours apart |
| Arowojolu et al. (2002) <sup>4</sup> | 573                   | 4                       | 1.5 mg as a single dose             |
| WHO (1998) <sup>5</sup>              | 976                   | 11                      | 0.75 mg in two doses 12 hours apart |
| Wu et al. (1999) <sup>6</sup>        | 643                   | 20                      | 0.75 mg in two doses 12 hours apart |
| Ho and Kwan (1993) <sup>7</sup>      | 410                   | 12                      | 0.75 mg in two doses 12 hours apart |
| Total                                | 5859                  | 98                      |                                     |

<sup>a</sup>Personal communication from Shangchun Wu, 16 February 2003. WHO, World Health Organization.



## Ectopic pregnancy following use of progestin-only ECPs

James Trussell, Allison Hedley and Elizabeth Raymond

*J Fam Plann Reprod Health Care* 2003 29: 249  
doi: 10.1783/147118903101197944

---

Updated information and services can be found at:  
<http://jfprhc.bmj.com/content/29/4/249.2.citation>

---

### Email alerting service

*These include:*

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

---

### Notes

---

To request permissions go to:  
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:  
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:  
<http://group.bmj.com/subscribe/>